

References

1. King N. Mild head injury: neuropathology, sequelae, measurement and recovery. *Br J Clin Psychol.* 1997;36 (Pt 2):161-184.
2. van der Naalt J. Prediction of outcome in mild to moderate head injury: a review. *J Clin Exp Neuropsychol.* 2001;23(6):837-851.
3. F K, LD C. *Epidemiology.* Washington DC: American Psychiatric Publishing; 2005.
4. RM R, C WJ. Myths and Mild Traumatic Brain Injury. *Psychological Injury and Law.* 2009;2(34).
5. Lange RT, Brickell TA, Ivins B, Vanderploeg RD, French LM. Variable, not always persistent, postconcussion symptoms after mild TBI in U.S. military service members: a five-year cross-sectional outcome study. *J Neurotrauma.* 2013;30(11):958-969.
6. Ponsford J, Willmott C, Rothwell A, et al. Factors influencing outcome following mild traumatic brain injury in adults. *J Int Neuropsychol Soc.* 2000;6(5):568-579.
7. Kashluba S, Paniak C, Casey JE. Persistent symptoms associated with factors identified by the WHO Task Force on Mild Traumatic Brain Injury. *Clin Neuropsychol.* 2008;22(2):195-208.
8. Luis CA, Vanderploeg RD, Curtiss G. Predictors of postconcussion symptom complex in community dwelling male veterans. *J Int Neuropsychol Soc.* 2003;9(7):1001-1015.
9. Losoi H, Silverberg ND, Wäljas M, et al. Recovery from Mild Traumatic Brain Injury in Previously Healthy Adults. *J Neurotrauma.* 2016;33(8):766-776.